

Measles outbreak investigation in a highly vaccinated community in the Centre region of Cameroon

Haddison C. Eposi,¹ Ngwafor A. Randolph,² Kagina M. Benjamin³

¹Saa District Health Service, Centre Regional Delegation of Public Health, Cameroon; ²Institute for Global Health, University College London, United Kingdom; ³Vaccines For Africa Initiative (VACFA), University of Cape Town, South Africa

Abstract

Background: Measles remains a threat in many African settings due to sub-optimal routine immunisation and catchup campaigns. The Global Vaccine Action Plan goal to eliminate measles by 2020 remains unmet as several countries reported an increase in cases in 2019. In Cameroon, a measles-rubella vaccination campaign was organised in 2019 to reduce the cohort of susceptible children. However, in 2020, eleven suspected cases of measles were notified in the Sa'a Health District and five were confirmed.

Objective: This report summarizes a measles outbreak investigation and contact tracing in a highly vaccinated community residing in the Sa'a Health District of Cameroon.

Methods: Outbreak investigations were carried out in the Sa'a, Nlong-Onambele and Nkolmgbana health areas from 18 to 21 February 2020. A register review from December 2019 to February 2020 was carried out in all health facilities of the affected health areas, followed by contact tracing in the community.

Results: Thirty households were visited in four neighbourhoods. Six missed Epidemiologically-linked cases were discovered in the community, bringing the total number of suspected and confirmed cases to 17. Thirty-five percent of the cases had not received any measles-containing vaccine; 35% of the cases were aged 5 years or older; 53% had history of travel. Community transmission only occurred in the Sa'a health area through a breakthrough case.

Conclusions: This outbreak investigation portrayed the role that adequate vaccination coverage plays in preventing widespread outbreaks. Nonetheless, community sensitisation and routine immunisation require strengthening in order to erase pockets of susceptible children.

Introduction

The global effort to fight measles and rubella have been guided by key recommendations and subsequent implementations of the 2012-2020 Global Strategic Plan for measles and rubella eradication, some of which are: effective surveillance; achieve and attain high levels of population immunity; outbreak preparedness and response.¹ However, the target set by the Global Vaccine Action Plan to eliminate measles in the World Health Organisation-African (WHO-AFRO) region by 2020 remains unmet.² In fact, according to the WHO, the AFRO region recorded a 700% increase in the number of measles cases in 2019.³ Cameroon is one of the countries in Africa that was reported to have a measles outbreak in 2019, with a total of 2729 cases, an 800% increase from 2018.⁴ One of the reasons for outbreaks of Vaccine-Preventable Diseases (VPDs) including measles, is sub-optimal vaccination coverages during routine immunisation. Therefore, measles catch up and mass campaigns are recommended in order to attain optimal population immunity and to mitigate the occurrence of outbreaks.¹

In Cameroon, like many other countries, measles vaccine is provided to children through routine immunisation services delivered by the Expanded Programme on Immunisation (EPI). The EPI in Cameroon targets 0-15 months old children. The measles immunisation programme began in Cameroon in 1965 and by 1974 the vaccine was administered only to 9-month-old children as a single dose.⁵ Since 2015 the vaccine is administered as a combined Measles-Rubella (MR) vaccine with first dose at 9 months of age. The national coverage for the first dose of the MR vaccine in 2019 was reported to be 71%.⁴ There were variations in coverage of the first dose for different regions. For example, three regions, including the Centre region recorded a coverage above the national target of 83%.⁴ The second dose of the MR vaccine targeting 15-months-old children was introduced into the EPI in 2020. As expected, uptake of the second dose in the Centre region remained very low after only 3 months of implementation. Uptake of the second dose is expected to improve with time as well as with mass vaccination campaigns.

Since 2002, Cameroon has carried out 5 follow up measles vaccination campaigns, with the last one taking place in 2015. Reports show that measles epidemics in the country peak every three years.⁵ The last

Correspondence: Haddison C. Eposi, Saa District Health Service, Centre Regional Delegation of Public Health, Cameroon.
Tel.: +237 699385785
E-mail: eposihaddison@yahoo.com

Key words: Measles; outbreak; investigation; vaccination; Cameroon.

Acknowledgements: Dr Tambasho Afizu, Central Technical Group-Expanded Programme on Immunisation, Ministry of Health, Cameroon; Mr Enyegue Mbia Guy, Sa'a District Health Service, Centre RDPH, Cameroon; Mr Okala Jean Didier, Nlong Onambele Integrated Health Centre, Sa'a DHS, Cameroon; Mme Ndzana Dior, Ayissile Integrated Health Centre, Sa'a DHS, Cameroon.

Contributions: HEC conceived the study and collected the data. HEC and NAR analysed the data and wrote the manuscript with supervision from KMB. All authors read and approved the final manuscript

Conflict of interest: The authors have no conflict of interest to declare.

Availability of data and materials: All data generated or analysed during this study are included in this published article

Ethics approval and consent to participate: The findings of this study were derived from routine immunisation activities so no ethical approval was required.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Received for publication: 11 December 2020.
Revision received: 23 March 2021.
Accepted for publication: 23 March 2021.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright: the Author(s), 2021
Licensee PAGEPress, Italy
Journal of Public Health in Africa 2021; 12:1775
doi:10.4081/jphia.2021.1775

peak was in 2015 thus the next follow up vaccination campaign was scheduled for 2018 in order to prevent an eventual peak. Unfortunately, the 2018 campaign was rescheduled to 2019. We speculate the rescheduling resulted to the resurgence in measles cases across the country due to an increase in the number of susceptible children.⁴ In effect, by the end of 2019, 47 of the 189 health districts in the country had notified 3 confirmed cases within 28 con-

secutive days, thus crossing the epidemic threshold.⁴ According to national guidelines, a response vaccination coverage targeting children 9-59 months is recommended in any locality where an epidemic occurs. It was therefore imperative that the measles campaign initially scheduled for 2018 be carried out, to reduce the number of susceptible children to prevent further outbreaks.

It is in this context that the 2019 national MR campaign coupled with Vitamin A supplementation and deworming targeting children aged 9 to 59 months was carried out in the Sa'a Health District (HD). The Sa'a HD is one of the 30 HDs of the Centre region of Cameroon. The MR campaign coordination was done at the District level with supervision from the Regional Technical Group-Expanded Programme on Immunisation. The co-ordination was similar to that of the Polio campaign done in 2018 as previously described by Haddison *et al.*⁶ The MR vaccine (Serum Institute, India) in vials of 10 doses was used for the campaign. The campaign took place from the 4th to the 10th of December 2019. Only the fixed post strategy in health facilities and temporary fixed post strategy in areas such as markets, churches and motor parks were used since this was an injectable vaccine. A total of 9164 out of 9308 children were vaccinated, resulting to an administrative campaign coverage of 97%.

To ascertain how successful the campaign was, a MR campaign coverage survey in the district was evaluated using WHO's cluster immunisation methodology.⁷ Three health areas-Sa'a, Nlong-Onambele and Nkolmbana were purposively chosen, with twenty-four clusters randomly chosen by balloting. Vaccination was ascertained either by visual inspection of the vaccination card distributed during the campaign or by recall. Out of the 479 children surveyed, 463 were vaccinated giving a MR campaign coverage of 96%.

Between the first and sixth epidemiological week of 2020, eleven suspected cases of measles were notified in the Sa'a HD where a successful MR campaign had been ascertained. Serum specimens from the suspected cases were sent to Centre Pasteur du Cameroun for analysis. Of the eleven, five specimens tested positive, surpassing the threshold for a measles epidemic. Therefore, sampling of suspected cases was suspended according to national policies pending an outbreak investigation and response. It is worth noting that there were zero confirmed measles cases in the district out of the 11 suspected cases notified in 2018 and 13 cases notified in 2019. Surprisingly, measles outbreaks were

declared in 15 of the 30 health districts of the Centre region within the same period. We report a measles outbreak investigation and contact tracing in a highly vaccinated community residing in Nlong-Onambele, Sa'a, and Nkolmbana health areas in the Sa'a HD.

Materials and Methods

Ethical considerations

The findings of this study were derived from routine immunisation activities so no ethical approval was required.

Setting

Investigations were carried out in three health areas; i) Sa'a HA which is the largest HA of the district and most populous with a population of 29307 inhabitants. This HA has the highest concentration of health facilities in the district including the district hospital; ii) Nlong-Onambele HA with a population of 6230 inhabitants served by one health facility; iii) and Nkolmbana HA, the smallest HA of the district with a population of 2274 inhabitants served by one health facility. This HA shares borders with Sa'a HA.

Investigation process

Investigations were carried out in two phases from 18 to 21 February 2020 (epidemiological week 8). Firstly, a register review from December 2019 to February 2020 was carried out in all health facilities of the affected HAs. Secondly, contact tracing was carried out in the community. The homes of the positive cases were visited first, then homes within a 100m radius of the homes of the positive cases and any suspected case found in the community were visited.

Case definitions

Suspected case: Anyone with fever and maculo-papular rash plus either cough, coryza or conjunctivitis.⁸

Confirmed case: Any suspected case with laboratory positive results for IgM antibody.⁸

Missed case: Any suspected case who was not notified but later identified in the community.

Naïve case: Any case who had never been vaccinated with a measles containing vaccine (MCV).

Breakthrough case: Any case who had received a dose of MCV.

Data collection and analysis

Data collected included socio-demo-

graphic characteristics of the cases, date of onset of the symptoms, date of consultation, vaccination and travel history as well as contact with other suspected cases. Data were subsequently entered and analysed using Microsoft Excel 2016 (Microsoft Corp., Redmond, Washington, United States). Proportions were calculated based on the total number of cases (confirmed, missed, suspected).

Results

A total of 30 households were visited in four neighbourhoods including the homes of the five positive cases. Majority of the cases from all the health facilities that were recorded in the register review were identified. Six missed epidemiologically linked cases were discovered in the community thus bringing the total number of suspected and confirmed cases to 17 from epidemiological week 1 to 8 (Figure 1).

Clinical presentation

The ages of the cases ranged from 10 months to 42 years (Table 1). All the cases presented with a fever and rash and were considered as having mild to moderate disease except for one case who developed rectorrhagia thus considered as severe disease (Table 2). No deaths were recorded.

Profiling of the cases

There was a similar proportion of breakthrough and naïve infections. Vaccination status could not be ascertained in 29.4% of the cases. Only one child between the ages of 9 months to 5 years had received two doses of the MR vaccine. Most of the cases (52.9%) had recently travelled out of the district (Table 1).

Investigation and contact tracing

Investigations began in the Ebogo neighbourhood in Saa HA where the confirmed case had been notified then moved to Nlong Onambele and Nkolmbana health areas. At the end of the investigations, the following facts were reconstituted based on the onset of symptoms in each health area.

During epidemiological week 1, the index case, a resident of Djoungolo HD (80km from Sa'a HD) visited her cousin, patient 2 who resided in the Bami neighbourhood in Saa HA. During the visit the index case developed symptoms of measles, which led to a cascade of transmissions in Saa HA from epidemiological week 1 to 5 (Figure 2). During epidemiological week 3, patient 2 developed symptoms of measles and was hospitalised in Ebogo where he also schooled. Investigators found out that

patients 3 and 4 (a missed case) who developed symptoms during epidemiological week 5 were his classmates. No other pupils were reported sick by the school principal. Back in the Bami neighbourhood investigators discovered patients 5 and 6 who were neighbours to patient 2.

Though patient 7 and his father (patient 8) had both consulted for measles in Ebogo during epidemiological week 5, investigators found no epidemiological link between them and patients 1-6. Investigators learnt that patients 7 and 8, residents of Djoungolo HD had come to Sa'a for the burial of a family member who allegedly died of measles in Djoungolo HD. Upon arrival they already had symptoms of measles. Investigators were unable to locate patients 9 and 10 who had been notified by the health centre.

In the Nlong Onambele HA, investigators found out that both cases notified by the health facility, patient 11 and patient 12, residents of Djoungolo HD had arrived Sa'a sick during epidemiological weeks 1 and 2

respectively in search of traditional medicine. No other cases were found in the community. Also, no epidemiological link could be found between them and the other cases in the district.

In the Nkolmbgana HA, during epidemiological week 3 patient 13 developed mild symptoms (fever) that were not classified as measles related until a week later. Investigators learnt patient 13 had visited




Figure 1. Epidemiological curve for measles cases from epidemiological week 1 to 8 by date of onset of symptoms.

Table 1. Characteristics of the measles cases.

Patient	Age	Notified	IgM sera	Onset of symptoms (epidemiological week)	Doses of MR	History of travel	Contact with another case	Health area of notification in Sa'a HD	Health district of residence
1	1.5y	No	ND	1	0	Yes	Yes	Sa'a	Djoungolo
2	5y	Yes	Pos	3	2	No	Yes	Sa'a	Sa'a
3	8y	Yes	Pos	5	0	No	Yes	Sa'a	Sa'a
4	6y	No	ND	5	1	No	Yes	Sa'a	Sa'a
5	4y	Yes	Neg	5	1	No	Yes	Sa'a	Sa'a
6	1.5y	Yes	Neg	5	0	No	Yes	Sa'a	Sa'a
7	2y	Yes	Pos	5	Unknown	Yes	Yes	Sa'a	Djoungolo
8	42y	Yes	Neg	5	Unknown	Yes	Yes	Sa'a	Djoungolo
9	14 mths	No	ND	5	0	No	Unknown	Sa'a	Sa'a
10	5y	Yes	Neg	6	1	No	Unknown	Sa'a	Sa'a
11	1.5 y	Yes	Neg	1	Unknown	Yes	No	Nlong Onambele	Djoungolo
12	20y	Yes	Pos	2	Unknown	Yes	No	Nlong Onambele	Djoungolo
13	10 mths	Yes	Pos	3	1	Yes	No	Nkolmbgana	Sa'a
14	1y	No	ND	2	1	No	Yes	Nkolmbgana	Sa'a
15	3y	No	ND	2	0	Yes	Yes	Nkolmbgana	Sa'a
16	1y	No	ND	2	0	Yes	Yes	Nkolmbgana	Sa'a
17	5y	Yes	Neg	6	1	Yes	No	Lepopomo	Sa'a

ND: Not done, Pos: Positive, Neg: Negative, y: years, mths: months.

Table 2. Distribution of symptoms among cases.

Symptoms	Naïve cases N = 6 (35.3%)	Breakthrough cases N = 6 (35.3%)	Indeterminate cases N = 5 (29.4%)
Fever	6	6	5
Rash	6	6	5
Cough	3	3	1
Conjunctivitis	2	3	1
Coryza	4	3	1
Rectorrhagia	0	1	0

measles in most settings.^{17,18} In Cameroon, measles is still endemic so in the event that these adults were vaccinated they would have benefited from this natural boosting. We can only speculate that they were naïve cases and there may be several more in the community.

During the period that the measles outbreak was declared in the Sa'a HD, other neighbouring urban HDs like Djoungolo had already declared measles epidemics. Many of the positive cases declared in Sa'a HD were residents of these districts or had recently travelled from these districts and arrived Sa'a while already infected. In many African settings, patients tend to leave the cities to rural areas in search of traditional or alternative treatment.¹⁹ This favours the importation of diseases which may lead to outbreaks in susceptible populations. This was the case in our setting and we assume the outbreak was very localised due to herd immunity.

Studies reporting outbreaks in vaccinated populations rarely describe onward transmission from breakthrough cases.²⁰⁻²² However, we observed this phenomenon in our setting. Patient 2 who had received 2 doses of the MR vaccine was the only contact to his classmates and neighbours in Sa'a health area. Moreover, he developed a severe form of the disease which is not in accordance with reports which describe mild disease in breakthrough cases.²⁰⁻²² Nonetheless, a similar case was described by Rosen *et al.*, in the United States.²³

Limitations

This outbreak investigation had several limitations. Firstly, cases were considered as breakthrough cases based on vaccination history alone instead of IgG serology. No IgG serology could be requested as the reference lab only conducts IgM serology for measles diagnosis. Also, vaccination history was based on recall, which may be biased. Secondly, since the community was highly vaccinated, some cases may have been missed if they presented with milder symptoms. Nonetheless, we believe there was no other community transmission because no cases were declared from epidemiological week 7. Lastly, this investigation was only carried out in three health areas so careful generalisation of the results is needed.

Conclusions

This outbreak investigation portrayed the role adequate vaccination coverage plays in preventing widespread outbreaks. Nonetheless, community sensitisation and routine immunisation requires strengthen-

ing in order to erase pockets of susceptible children.

References

1. WHO. Global Measles and Rubella strategic plan 2012-2020. Accessed 04 June 2020. Available from: https://www.who.int/immunization/diseases/measles/global_coordination/en/index3.html
2. WHO. Global Vaccine Action Plan 2011-2020. Accessed 04 June 2020. Available from: https://www.who.int/immunization/global_vaccine_action_plan/GVAP_doc_2011_2020/en
3. World Health Organisation. New measles surveillance data for 2019. Accessed 04 June 2020. Available from: <https://www.who.int/immunization/newroom/measles-data-2019/en/>
4. Ministry Of Health. Expanded programme on immunisation annual report, 2019. Cameroon: Ministry Of Health.
5. Cummings DAT, Moss WJ, Long K, et al. Improved measles surveillance in Cameroon reveals two major dynamic patterns of incidence. *Int J Infect Dis* 2006;10:148-55.
6. Haddison EC, Ngono D, Kouamen GT, Kagina BM. Successful polio supplementary immunisation activities in a security compromised zone - Experiences from the Southwest region of Cameroon. *Vaccine* 2018;36:6961-7.
7. World Health Organization Vaccination Coverage Cluster Surveys: Reference Manual. 2018 Accessed 09 January 2020. Available from: https://www.who.int/immunization/documents/who_ivb_18.09/en/
8. World Health Organisation. WHO-recommended surveillance standard of measles. Accessed 10 June 2020. Available from: https://www.who.int/immunization/monitoring_surveillance/burden/vpd/surveillance_type/active/measles_standards/en/
9. World Health Organisation. Measles vaccine: WHO position paper-April 2017. *Weekly Epidemiol Record* 2017;92:205-28.
10. Smith LE, Amlôt R, Weinman J, et al. A systematic review of factors affecting vaccine uptake in young children. *Vaccine* 2017;35:6059-69.
11. Vonasek BJ, Bajunirwe F, Jacobson LE, et al. Do maternal knowledge and attitudes towards childhood immunizations in rural Uganda correlate with complete childhood vaccination? *PLoS One* 2016;11:e0150131.
12. Salmon DA, Dudley MZ, Glanz JM, Omer SB. vaccine hesitancy: causes, consequences, and a call to action. *Am J Prev Med* 2015;49:S391-8.
13. Ashbaugh HR, Cherry JD, Hoff NA, et al. Measles antibody levels among vaccinated and unvaccinated children 6-59 months of age in the Democratic Republic of the Congo, 2013-2014. *Vaccine* 2020;38:2258-65.
14. Carazo S, Billard MN, Boutin A, De Serres G. Effect of age at vaccination on the measles vaccine effectiveness and immunogenicity: systematic review and meta-analysis. *BMC Infect Dis* 2020;20:251.
15. Dine MS, Hutchins SS, Thomas A, et al. Persistence of vaccine-induced antibody to measles 26-33 years after vaccination. *J Infect Dis* 2004;189:S123-30.
16. Ramsay ME, Moffatt D, O'Connor M. Measles vaccine: a 27-year follow-up. *Epidemiol Infect* 1994;112:409-12.
17. Whittle HC, Aaby P, Samb B, et al. Effect of subclinical infection on maintaining immunity against measles in vaccinated children in West Africa. *Lancet* 1999;353:98-102.
18. Paunio M, Hedman K, Davidkin I, et al. Secondary measles vaccine failures identified by measurement of IgG avidity: high occurrence among teenagers vaccinated at a young age. *Epidemiol Infect* 2000;124:263-71.
19. James PB, Wardle J, Steel A, Adams J. Traditional, complementary and alternative medicine use in Sub-Saharan Africa: a systematic review. *BMJ Glob Health* 2018;3:e000895.
20. Sundell N, Dotevall L, Sansone M, Andersson M, Lindh M, Wahlberg T, et al. Measles outbreak in Gothenburg urban area, Sweden, 2017 to 2018: low viral load in breakthrough infections. *Euro Surveill* 2019;24:1900114.
21. Avramovich E, Indenbaum V, Haber M, et al. Measles outbreak in a highly vaccinated population - Israel, July-August 2017. *MMWR Morb Mortal Wkly Rep* 2018;67:1186-8.
22. Berry L, Palmer T, Wells F, et al. Nosocomial outbreak of measles amongst a highly vaccinated population in an English hospital setting. *Infect Prev Practice* 2019;1:100018.
23. Rosen JB, Rota JS, Hickman CJ, et al. Outbreak of measles among persons with prior evidence of immunity, New York City, 2011. *Clin Infect Dis* 2014;58:1205-10.